

**REMOVAL OF CU (II) IONS FROM WASTEWATER BY USING DRIED
EUCALYPTUS LEAVES USING BATCH ADSORPTION AND STUDY THE
ADSORPTION THERMODYNAMICS**

IHAB . A. ALTAMEEMI*
RIHAB R. ABDUL-REZAQ**

*Department Of Chemistry, College Of Education for Pure Sciences, University Basra, Iraq

**Department Of Maths, College Of Education for Pure Sciences ,University Basra, Iraq

Abstract

A new simple method for the treatment of waste water effluent containing Cu(II), was developed using dried eucalyptus leaves a low-cost natural . Batch experiments were conducted to determine the effects of varying adsorbent weight, pH, contact time, metal ion concentration and temperature of adsorption. The adsorption of Cu (II) was found to be maximum(80%) at pH 8, at 25°C, metal ion concentration 100 ppm , contact time 60 min and Speed Shake(185rpm) . adsorption capacity of Cu (II) were found maximum (82.5%) at optimum conditions. Freundlich isotherm was found to be suitable for the adsorption of Cu(II) functional groups[(C-N), (C-O),(C=O),(O-H)] identification was given using FTIR spectrophotometry.

Keywords: adsorption, copper, Freundlich isotherm, Langmuir isotherm, dried eucalyptus leaves.

1.Introduction

The toxicity of heavy metals in the environment is still major concerns of human life; because they accumulate in living tissues throughout the food chain which has humans at its top. The danger of these heavy metals is due to poisoning, cancer, and brain damage[1].The treatment of water and waste water containing heavy metal is very demanding. Such methods of treatment include precipitation [2], flotation [3], biosorption [4][6], electrolytic recovery, membrane separation[7], removal by adsorption on minerals [8,9] and activated carbon adsorption [10,11]. Heavy metals are the most important pollutants in the environment and widely found on water surfaces from natural and industrial sources[11] . Among metals, the two important pollutant are Cu and Ni cations. Cu ions can deposit in brain, skin, liver, pancreas, and myocardium causing serious toxicological effects on humans[12, 13]. Cu (II) is a commonly toxic metal in natural ecosystems. The most common adverse health effect of nickel in humans is an allergic reaction in direct contact with the skin caused by jewellery or other items containing Cu. There are several methods for removing heavy metals from waste water such as adsorption, chemical precipitation, ion exchange, electrochemical treatment, solvent extraction, and membrane separation[14, 15].

Nowadays, a number of low cost adsorbent materials such as zeolites are available to remove metal ions[16-17-18].

1:1 EXPERIMENTAL SECTION

Adsorption experiments

For the adsorption experiment, the effect of adsorbent weight on the adsorption of the ions was investigated (0.5 -2g) of the adsorbent was weight respectively into conical flasks. 20 ml of 100 mg/L solution of each of the metal ions solution was added and the mixture shaken at 185 rpm for 60 min. After the biosorption experiment, the biosorbent was separated from the solution by filtering through a Whatman (0.45 µm) filter paper and the filtrate was analyzed for the Cu (II) ions concentration. The Cu (II) ions concentrations before and after the biosorption were determined by using Flame Atomic Absorption Spectrophotometer. (Shimadzu 6000). The equilibrium Cu(II) ions sorption capacity could be calculated as equation bellow;

$$Q_e = V(C_0 - C_e) / m \dots (1) [19]$$

Where ; Q_e is the equilibrium Cu (II) ions sorption capacity (mg/g), V is the suspension volume (l), m is the mass of durian material (g), C_0 = Initial concentration of solution, C_e = Concentration of the solution after adsorption. To determine the effect of concentration on the adsorption of the metal ions, 2g of the adsorbent, being the optimum adsorbent weight in the previous experiment was added to 50 ml each of varying concentrations(between 5 - 100 mg/L) of the metal ion solutions. The mixtures were shaken and the concentration of the metal ions adsorbed was determined. The effect of contact time was also investigated by adding 2g of the adsorbent to 20 ml of 50 mg/L of Cu (II) ions being their respective optimum adsorption concentrations; and shaking using varying contact times(30 –180 min) and the percentage of adsorbed ions determined. The effect of pH on adsorption of the metals was investigated using 0.5g of the adsorbent and 20 ml of 100 mg/L of Cu (II) ions. The mixture was shaken for 60 min for Cu (II) and the amount of ion adsorbed was determined[20].

Preparation of adsorbent

Eucalyptus leaves were collected from gardens of AL-basrah University, basrah, Iraq. The leaves were extensively washed with deionize water to remove dirt ,dried in an oven at 90°C° for a period of 1.5 hr, then ground and screened to obtain the average particle size 150µm.the powder

was preserved in glass bottles for use as adsorbent, The some physical properties of the chosen adsorbent are listed in Table 1.

Table 1

Properties	Value
Porosity(%)	82.99 ±0.349
Langmuir surface area(m ² /gm) 2	1.7656 m ² /g
BET Surface Area	1.1312 m ² /g
Pore Volume	9.709 cm ³ /g
Partical density	0.718 g/cm ³
Bulk density	0.688 gm/cm ³

2: RESULTS AND DISCUSSION

2:1. FT-IR Characterization of Eucalptus Leaves

FT-IR apparatus type shimadzu (4000-400 cm⁻¹ P) was carried out to identify the functional groups and structural in the Eucalptus leaf powder that might be involved in the adsorption process. FT-IR analysis was carried out in order to identify the functional groups in the Eucalptus leaf powder that might be involved in the adsorption process. The FTIR spectrum in the range of 400-4000cm is shown in Fig.1.as shown in the figure, the spectrum displays a number of adsorption peak, which indicates the complex natural of the material examined stretching, which is consistent with the peak at 1160.28 and 1340cm assigned to alcoholic C-O and C-N stretching vibration. The absorption band wave numbers of 2990-2870 cm can be assigned to CH and -CH₂ stretching, respectively [21].The absorption band wave number of ketone group is about 1690cm.that result of binding of this group with the cations of metal .Show table(2).

Table (2) explains the effective groups

Wavelength(cm ⁻¹)	Structural and Functional group	Functional group
1600-1690	C=O	ketone group
1700-1720	C=O	carboxylate groups
3000-3600	O-H	Hydroxyl group
2870-2990	C-H	Hydrocarbon
1310-1340	N-H	Amines
1050-1160	C-O	Carbon oxygen group




Fig (1) FT- IR Eucalyptus leaves powder

2:2.Effect of pH

The pH is one of the most important parameters of biosorption of heavy metals, The biosorption of Cu (II) by Eucalyptus erectus leaf powder at different pH values(3-10) is presented in Fig2. The optimal pH for removal of Cu (II) was(10). The removal percentages of, Cu (II) ion on Eucalyptus powder at an initial concentration of 100 ppm and at an initial pH=(10) ,using 0.5g of the adsorbent with 100mLof Cu (II) ions solution (80%).At pH higher than 10 both metals were precipitated due to the formation of hydroxides and removal due to sorption was very low .The minimal adsorption at low pH may be due to the higher concentration and high mobility of the H⁺, which are preferentially adsorbed rather than the metal ions . At higher pH values, the lower number of H⁺ and greater number of ligands with negatives charges results in greater Cu (II) adsorption. For example, carboxylic groups (-COOH) are important groups for metal uptake by biological materials . At pH higher than 3-4, carboxylic groups aredeprotonated and negatively charged. Consequently, the attraction of positively charged metal ions would be enhanced [22].




Fig (2) Effect of pH

2:3. Effect of adsorbent weight

One of the parameters that strongly affect the sorption capacity is weight of the adsorbents. With the fixed metal concentration it can easily be inferred that the percent removal of metal ions increases with increasing weight of the adsorbents as shown from Fig. 3 This is due to the greater availability of the exchangeable sites or surface area at higher concentration of the adsorbent[23]. The removal percentages of Cu(II) ion Eucalyptus powder at an initial concentration of 100 ppm and at an initial pH=(8), using at different weight of the adsorbents(0.5_3g) with 50mL of Cu (II) ions solutions(96.9%).




Fig (3) Effect of adsorbent weight

2:4. Effect of metal ions concentrations

The removable efficiency decreases with increasing initial concentrations from 10-100 mg L⁻¹, shown in Figure 4. Maximum adsorption obtained at minimum concentration of metal ions. The different lead solutions with concentration ranging from 10mg/l to 100 mg/l were agitated at 250 rpm for 80 minutes with adsorbent dose of 0.5 gm,. The percentage adsorption of Cu(II) decreases as the initial concentration of increased from 10 mg/l to 100 mg/l. Maximum adsorption obtained at minimum concentration of metal ions. Such behavior can be because of the unchanging number of available active sites on the adsorbent here the amount of adsorbent was constant[24].




Fig (4) Effect of Initial Concentration of Pb(II)ions

2:5.Effect of contact time

The effect of contact time on the biosorption of lead and copper ions onto nano sawdust is studied at different times ranged from 30-180 min. and optimum pH Cu(II)]. Figure 5 shows that the removal of Cu(II) as a function of contact time, where the removal percentage of the two metal ions increase rapidly with increasing contact time (i.e., high adsorption rate). For lead, minimum percentage removal was 80.8% at time of 30 min and it increased to maximum value 82.5% at time of 60 min, whereas, for copper, minimum value was 82.6% at time of 90 min to maximum value of 83% at time of 120 min. After maximal time, increasing in contact time is not effective in the adsorption process.[25] The rate of the biosorption of the two metal ions onto the surface of the nano sawdust occurred in two steps. The biosorption was very fast initially as a large number of vacant surface sites are available for binding with metal ions. As the surface adsorption sites become exhausted, the uptake rate is controlled by the rate at which the adsorbate is transported from the exterior to the interior sites of the adsorbent particles.[26] Then biosorption reached equilibrium as a result of the reduction of available sites which are difficult to be occupied due to repulsive forces between metal ions biosorbed onto the biosorbent surface and the bulk phase [27].




Fig (5) Effect of contact time

2:6. Effect of temperature on the adsorption rate

The Increase of temperature from 30 to 60 C° increased the adsorption of Cu (II) ions indicating the process to be endothermic. The increases in uptake of Cu (II) ions with temperature may be due to the desolvation of the adsorbing species, the changes in the size of pores, and the enhanced rate of intraparticle diffusion of adsorbate, as diffusion is an exothermic process. The biosorption was found to increase with increase in temperature at (30-55P OPC)and the sorption capacity (Q) was also found to increase. The interactions are

found to be exothermic in nature [28],[29] for which the evaluation of thermodynamic. This is shown in Figure(6). The removal percentages of Cu (II)ions on Eucalptus powder was at different temperature from(30-60) and at an initial pH=8,using(0.5g of adsorbent)with 20mL of Cu (II)ions solutions (91.9%.) [30].




Fig (6) Effect of temperature on the adsorption rate

3. Isotherms

The adsorption isotherm experiment was carried out by mixing 1 gram of adsorbent and 100 mL of metal ion solution of various concentration; 20-100 mg/L with pH 8, shaking at 150 rpm about an hour for reaching equilibrium. Then the remaining concentration of metal ion in solutions were examined using AAS for Langmuir adsorption isotherm model and Freundlich adsorption isotherm model.

Langmuir model:

$$q_e = \frac{q_m K_L C_e}{(1 + K_L C_e)}$$

and can be linearized to be

$$\frac{C_e}{q_e} = \frac{C_e}{q_m} + \frac{1}{K_L q_m} \dots\dots\dots(2)$$

Freundlich model:

$$q_e = K_F C_e^{\frac{1}{n}}$$

which can be written in the linear form as

$$\log q_e = \log K_F + \frac{1}{n} \log C_e \dots\dots\dots(3)$$

where C_e is equilibrium concentration (mg/L), q_e is adsorption capacity (mg/g), q_m is maximum adsorption capacity (mg/g), K_L is Langmuir adsorption constant and K_F is Freundlich adsorption constant. The plot of C_e/q_e versus C_e for Langmuir's adsorption model will give the straight line with slope of $1/q_m$ and intercept of $1/(K_L q_m)$. For Freundlich's adsorption model, plot of $\log q_e$ against $\log C_e$ also will be linear relationship. The relative coefficients of these models were calculated using linear least-squares fitting [31]. Since the relationship between the amount of substance adsorbed per unit mass of adsorbent at constant temperature and its concentration in equilibrium, adsorption isotherm, is very important in determining the adsorption capacity of metal ions onto the adsorbent. Adsorption isotherm describes the interaction between adsorbate and adsorbent materials. The experimental results were fitted to the equations according to Langmuir adsorption model and Freundlich model as Eq(2) and Eq(3), respectively. And the experimental data and calculated values are in Table 3. As shown in Figure (7) and regression coefficient in Table1, the experimental data fitted to Langmuir's adsorption model with R^2 around 0.58 and also fitted to Freundlich's adsorption model with R^2 around 0.95 as shown in Figure 4 for Cu (II) [32] .

Table (3)

Ion	Langmuir adsorption model			Freundlich adsorption model		
	a (mg/g)	b (L/g)	R^2	n	K_f	R^2
Cu	0.0538	0.0305-	0.88	0.04	0.736	0.96




Fig (7) Isotherms

4. Thermodynamic Study

The consideration of energy and entropy in any adsorption process is important for determining whether the process will be spontaneous or not. As a result, thermodynamic parameters are significant for the application of this material and process to real life scenarios. Equations (4), (5) and (6) were used to calculate the change in free energy ΔG° , change in enthalpy ΔH° , and change in entropy ΔS° which:

$$\Delta G^\circ = -RT \ln K \dots\dots\dots(4)$$

$$\log K = (\Delta S^\circ / 2.303R) - (\Delta H^\circ / 2.303 RT) \dots\dots\dots (5)$$

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ \dots\dots\dots (6)$$

As K is the equilibrium constant, T is the absolute temperature (K°) and R is the gas constant (33).

Van,t Hoff equation was used to estimate the values of ΔH° and ΔS° from the intercept, and slope of the plot which was $\ln K$ vs. $1/T$ (fig. 8). Table (4) showed the thermodynamic parameters values for the adsorption processes. From this the table, adsorption was found exothermic (ΔH° was negative), reaction was spontaneous (ΔG° was negative), and solid solution interface was random because ΔS° was increased. The change in adsorption enthalpy was measured and found -20 to 40 KJ.mol⁻¹, compared to chemisorption which was -400 and -80 KJ.mol⁻¹. values of ΔH° physisorption was found the dominant mechanism(34,35).




Fig (8) Effect of temperature on the adsorption rate.

Table (4)

heavy metal	Temperature K°	ΔG° KJ.mole-1	ΔH° KJ.mole-1	ΔS° J.mole-1.K-1
Cu	303	-15.41	-9.73	18.75
	313	-15.59		18.71
	323	-15.78		18.72
	333	-15.97		18.747

REFERENCES

1. Al-Garni SM Biosorption of lead by Gram-ve capsulated and non-capsulated bacteria. Water SA., 31,345–50,(2005).(
2. O.J. Esalah, M.E. Weber and J.H. Vera, “Removal of lead, cadmium and zinc from aqueous solutions by precipitation with sodium di-(n-octyl) phosphinate” Can. J. Chem. Eng., 78, 948–954,(2000)
3. A.I. Zouboulis, K.A. Matis, B.G. Lanara and C.L. Neskovic,” Removal of cadmium from dilute solutions by hydroxy apatite. II. Floatation studies” ,Sep. Sci. Technol., 32, 1755–1767,(1997).
4. Y.S. Ho, J.C.Y. Ng and G. McKay, “Removal of lead (II) from effluents by sorption on peat using second-order kinetics”, Sep. Sci. Technol., 36, 241–261, (2001).(
5. C. Hall, D.S. Wales and M.A. Keane, Copper removal from aqueous systems: biosorption by pseudo monassyringae, Sep. Sci. Technol., 36,223–240,(2001).(
6. Y. Sag, B. Akcael and T. Kutsal, Ternary biosorptionequilibria of chromium (VI), copper (II) and cadmium (II) onRhizopusArrhizus, Sep. Sci. Technol.,37, 279–309,(2002).(
7. L. Canet, M. Ilpide and P. Seat, Efficient facilitated transport of lead, cadmium, zinc and silver across a flat sheet-supported liquid membrane mediaed by lasalocid A, Sep. Sci. Technol., 37, 1851–1860,(2002).(
8. S. Ahmed, N. Khalid and M. Dand, Adsorption studies of lead minerals from aqueous media, Sep. Sci. Technol., 37, 343–362, (2002).(
9. D.B. Weirich, R. Hari, P. Behra and L. Sigg,Adsorption of Cu, Cd and Ni on goethite in presence of natural ground water ligands, Environ. Sci.Technol., 36, 328–336,(2002).(
10. V. Ravindran, M.R. Stevens, B.N. Badrihya and M. Pirbazari, Modeling the sorption of toxic metals on chelant-impregnated adsorbent, AICHE J., 45,1135–1146,(1999).(
11. Wang S, Wu H (2006) Environmental-benign utilisation of fly ash as low-cost adsorbents. J Hazard Mater.501–482 ,136
12. Wang S, Terdkiatburana T, Tade MO (2008) Single and co-adsorption of heavy metals and humic acid on fly ash. SeparPurif Tech 58, 353–8.
13. Veli S, Alyüz B (2007) Adsorption of copper and zinc from aqueous solutions by using natural clay. J Hazard Mater 149, 226–33.
14. Huang CP, Blankenship DW (1984) The removal of mercury(II) from dilute aqueous solution by activated carbon. Water Res 18, 37–46.
15. Panday KK, Parsad G, Singh VN (1985) Copper(II(removal from aqueous solutions by fly ash. Water Res.73–869 ,19
16. Crini G (2006) Non-conventional low-cost adsorbents for dye removal: A review. Bioresour Tech 97 , .85–1061
17. Cochrane EL, Lu S, Gibb SW, Villaescusa I (2006) A comparison of low-cost biosorbents and commercial sorbents for the removal of copper from aqueous media. J Hazard Mater 137, 198–206.
18. Li XG, Feng H, Huang MR (2008) Natural sorbents for mercury-ion removal. ProgrChem 20, 233–8.
19. S.E.Elaigwu, L.AUzman, G.V.Awolola, G.B.Adebayo, R.M.K.Ajayi: Adv.in. Appl.Sci, (2009) ,3(3),. 37,442-446.

20. Removal of pb (II), cd (II) Ions onto dried Conocarpus erectus leaves using batch adsorption and study the adsorption thermodynamics Ihab. A. Al-tameemi¹, Tarikak. Nasser² and Thuraya. M. Al Department of Chemistry, College of Education, University of Basrah, Iraq, Department of Marine Chemistry, Basra Marine Center, Basra University, Iraq, J. Chem. Pharm. Res., 2012, 4(12):4961-4968.
21. Pretsch, E., Clem, T., Seihl, J., and Simon, W. Spectra Data for Structure Determination of organic Compounds. Berlin: Springer-Verlag., (1981).
22. Ihab.A.altameemi^{1*} "removal of cadmium(ii) onto dried eucalyptus erectus leaves using batch adsorption and study the adsorption thermodynamics" Dept. of Chemistry, College of Education for Pure Sciences, University Basra, Iraq Impact Factor 3.114, ISSN: 2320-5083, Volume 5, Issue 8, September 2017
23. [S.H. Lee, C.H. Jung, H. Chung, M.Y. Lee, J.-W. Yang, Removal of heavy metals from aqueous solution by apple residues, Process. Biochem. 33, 205–211, (1998).
24. A low cost material, banana peel for the removal of lead (ii) from aqueous solutions Priyanka Kumari Volume: 04 Issue: 06 | June -2017 Department of Chemistry, University of Delhi, Delhi-110007, India.
25. International Journal of Latest Technology in Engineering, Management & Applied Science (IJLTEMAS) Ocimum Sanctum Sreelakshmi C.D Malabar College of Engineering and Technology, Kerala Technological University www.ijltemas.in Volume VI, Issue IV, April 2017 | ISSN 2278-2540 Page 85
26. Malekpour, A.; Khodadadi, M. Albumin-functionalized magnetic nanoparticles as an efficient sorbent for removal of Pb(II), Cd(II), Cu(II) and Cr(II) ions from aqueous solutions. RSC Adv. 2016, 6, 14705–14711
27. Modified Sunflower Seed Husks for Metal Ions Removal from Wastewater Siriwan Srisorrachtr Department of Chemical Engineering, Faculty of Engineering, Srinakharinwirot University, Nakhon Nayok 26120, Thailand. siriwans@g.swu.ac.th VOL. 57, 2017.
28. A new simple method for the treatment of waste water containing Cu (II) and Zn(II) Ions using adsorption on dried Conocarpus erectus leaves. Ihab.A.altameemi¹, Thuraya.M.AP and Tarikak. Nasser Department of Chemistry, College of education for pure sciences, University Basra, Iraq. Marine chemistry-Basra marine center-Basra university, Iraq .Received P6-11-2012, Accepted 12-3-2013
29. Khan, T.A.; Dahiya, S.; Ali, I. Use of kaolinite as adsorbent: Equilibrium, dynamics and thermodynamic studies on the adsorption of rhodamine B from aqueous solution. Appl. Clay Sci. 2012, 69, 58–66.
30. Anees Ahmad, Mohd. Rafatullah, Othman Sulaiman, Mahamad Hakimi Ibrahim, Yap Yee Chii, Bazlul Mobin Siddique "Removal of Cu(II) and Pb(II) ions from aqueous solutions by adsorption on sawdust of Meranti wood" Desalination 250, 300–310 (2009)
31. Lavecchia R., Medici F., Patterer M.S., Zuerro A., 2016, Lead removal from water by adsorption on spent coffee grounds, Chemical Engineering Transactions, 47, 295-300, DOI: 10.3303/CET1647050.
32. Madhava, M., Ramana, K., Seshaiyah, K., Wang, M., Chang, S.W., 2009, Removal of some metal ions by activated carbon prepared from Phaseolus aureus hulls, Journal of Hazardous Materials, 166, 1006-1013.
33. Huang, Ruihua; Wang, Bo; Yang, Bingchao; Zheng, Dongsheng and Zhang, Zengqiang, j. Desalination, (2011), Vol.280, pp.297-304.
34. Modification of multi-wall carbon nanotubes for the removal of cadmium, lead and arsenic from wastewater. Veličković, Z.S.; Bajić, Z.J.; Ristić, M.D.; Djokić, V.R.; Marinković, A.D.; Uskoković, P.S.; Vuruna, M.M Digest J. Nanomater. Biostruct. 2013, 8, 501–511.
35. Huang, Ruihua; Wang, Bo; Yang, Bingchao; Zheng, Dongsheng and Zhang, Zengqiang, j. Desalination, (2011), Vol.280, pp.297-304.